

RAPID H α VARIABILITY IN T CORONAE BOREALIS *

R.ZAMANOV^{1,2}, A. GOMBOC^{1,3}, M. F. BODE¹, J.M. PORTER¹, AND N.A. TOMOV²

¹Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Birkenhead, CH41 1LD, UK

²National Astronomical Observatory Rozhen, POB 136, Smoljan, BG-4700, Bulgaria

³Faculty of Mathematics and Physics, University in Ljubljana, Jadranska 19, 1000 Ljubljana, Slovenia

to be submitted to PASP

ABSTRACT

We report our search for variability in the H α emission line of the recurrent novae T CrB with time resolution 10-15 minutes. This is comparable with the time scale of the photometric flickering observed in this object. This is the first time that observations of the short time scale variation in emission lines have been made in this object. On two nights (990106 and 990107) we detected statistically significant variability (at 0.99 confidence level) in the line profile of H α . This variability is confined in the central part of the emission line ($\pm 100 \text{ km s}^{-1}$) although FWZI(H α) is $\sim 800 \text{ km s}^{-1}$. The variability in the line profile is accompanied with variability of the total equivalent width, EW(H α), $\pm 8\%$ for 990106, and $\pm 6\%$ for 990107 (calculated from the mean value of EW). Assuming Keplerian motion, the variability is generated at a distance $\sim 20\text{-}30 R_{\odot}$ from the white dwarf, which is approximately the radius of the ring the stream of gas forms as it flows away from L₁. On three other nights we are only able to put upper limits on the variability, $\Delta \text{EW}(\text{H}\alpha) \pm 2\%$ for 980415, $\pm 4\%$ for 980802, and $\pm 3\%$ for 980803.

Subject headings: stars:individual: T CrB – binaries: symbiotic – binaries:novae, cataclysmic variables

1. INTRODUCTION

The short term photometric behaviour of symbiotic stars has been investigated in many systems (e.g. Dobrzycka et al., 1996; Sokoloski et al., 2001). However, there are no systematic searches for rapid *spectral* variations. Until now the search for rapid spectral variability has been undertaken in only three symbiotics: CH Cyg, MWC 560, and RS Oph. All three stars are known to exhibit flickering and have collimated outflows. Rapid spectral variability in the Balmer lines of CH Cyg on time scales of ~ 1 hour have been detected and is probably connected with blobs ejected from a white dwarf, acting as a propeller (Tomov et al. 1996). For RS Oph, Sokoloski (2002) reported variability on time scales of hours in the HeII λ 4686 line although no variability in H β was detected. No significant variability has been detected in MWC 560 (Tomov et al. 1995).

In a few cataclysmic variables (CVs), which are closely related to symbiotic stars, rapid spectral changes are visible in optical and UV lines as a result of a variable accretion disk wind: BZ Cam (Ringwald & Naylor 1998), V592 Cas (Witherick et al. 2003), RW Sex (Prinja et al. 2003), V 603 Aql (Prinja et al. 2000a), BZ Cam (Prinja et al. 2000b).

The detection of variability on time scales of tens of minutes in symbiotic stars and cataclysmic variables provides the motivation for this study. T CrB consists of a red giant and a hot component, most probably a white dwarf. T CrB can be classified as a cataclysmic variable, symbiotic star, and recurrent nova and accretes at a rate $1 \times 10^{-8} M_{\odot} \text{ yr}^{-1}$ (Selvelli et al. 1992). This places it among the cataclysmic variables with the highest accretion rates (see Warner 1995, p.476) and for such high ac-

cretion rates an accretion disk wind could be expected. Here we report our search for signatures of variability in time-resolved optical spectroscopy in the H α line of T CrB, with time resolution $\sim 10\text{-}15$ minutes. This resolution is comparable with the time scale of flickering (see for example Zamanov et al. 2004).

2. OBSERVATIONS

The observations were performed with the Coudé spectrograph of the 2.0 m RCC telescope at the Bulgarian National Astronomical Observatory ‘‘Rozhen’’. The spectra have a dispersion of $0.2 \text{ \AA pixel}^{-1}$. The normal CCD reduction procedures (i.e. bias subtraction, flat fielding, wavelength calibration, etc) have been undertaken in the IRAF environment. A journal of observations is given in Table 1. The signal to noise ratio (SNR) achieved on the individual exposures is 35-65.

The averaged spectra, normalised to the local continuum, are presented in Fig.1, along with a comparison spectrum of the red giant HD135530.

3. RESULTS

3.1. Average, fractional and temporal variance spectra

The prominent emission line in Fig. 1 is H α . On a time scale of years the intensity of the H α line varies from $\text{EW}=0.5 \text{ \AA}$ up to $35 \text{ \AA}$ (Iijima 1990, Anupama & Prabhu 1991, Zamanov & Marti 2000, Stanishev et al. 2004). During the search for rapid variability we have observed a range of different profiles. An atlas of the different H α profiles is given in Stanishev et al. (2004). In the spectra obtained on 15th April 1998 the line is single peaked, and the central absorption dip is not visible. During August 1998, the profile is a little unusual, with 3 peaks or 2 absorption dips. During January 1999, the profile exhibits double peaks with a well defined central dip, which is partly due to the H α absorption line of the red giant.

* based on observations obtained in National Astronomical Observatory Rozhen, Bulgaria

³ e-mail: rz@astro.livjm.ac.uk; ag@astro.livjm.ac.uk; mfb@astro.livjm.ac.uk; jmp@astro.livjm.ac.uk

TABLE 1

H α OBSERVATIONS OF T CrB. IN THE TABLE ARE GIVEN THE DATE OF OBSERVATIONS, THE BEGINNING AND THE END OF SPECTROSCOPIC OBSERVATIONS AND TRUNCATED JULIAN DAY. ORBITAL PHASE IS CALCULATED USING THE EPHEMERIS OF FEKEL ET AL. (2000). THE FOLLOWING COLUMNS GIVE WAVELENGTH COVERAGE, NUMBER OF EXPOSURES IN EACH NIGHT AND THE EXPOSURE TIME OF THE INDIVIDUAL EXPOSURES. FOR THE EW(H α) WE GIVE THE MEAN, STANDARD DEVIATION, PLUS THE SMALLEST AND THE LARGEST VALUES FOR EACH NIGHT.

Date	UT _{start} - UT _{end}	TJD	phase	wavelength	N x Exp.time	EW(H α)	EW _{min} - EW _{max}
yymmdd	hh:mm - hh:mm	mid		coverage		[Å]	[Å]
				[Å]			
980415	22:37 - 01:43	50919.512	0.187	6510 - 6620	10 x 15 min	20.84 ± 0.31	20.39 - 21.36
980802	18:50 - 20:58	51028.328	0.665	6510 - 6620	12 x 10 min	13.39 ± 0.23	12.72 - 13.81
980803	18:23 - 21:03	51029.305	0.669	6510 - 6620	15 x 10 min	13.97 ± 0.25	13.63 - 14.45
990106	02:59 - 04:13	51185.652	0.356	6500 - 6700	7 x 10 min	9.77 ± 0.62	8.90 - 10.51
990107	01:50 - 02:51	51186.598	0.360	6507 - 6707	3 x 20 min	7.92 ± 0.45	7.45 - 8.36




FIG. 1.— The nightly average spectra for all five nights. A spectrum of the red giant HD135530 (M2III) is shown for comparison at the bottom.

The total EW(H α) was measured on each exposure in our sample. The results are presented in Fig. 2. In the first three nights the variability is less than three times the estimated errors. On 990106 and 990107 it is about 5 times the error of the individual measurements.

The measurements of the EW depend strongly on the continuum placement. The continuum was defined as a straight line or spline fitting, using parts of the spectrum located on both sides of H α . We used different wavelength intervals to produce different normalisations. This leads to changes of the absolute value of EW, but keeps the relative distributions of the points the same. The errors of the EW have been calculated from four measurements, with different normalizations.

In order to search for the variability in the line profile of H α line during an individual night, we calculate the fractional variance spectrum based on all the data in each




FIG. 2.— Measurements of the EW(H α). The vertical error bars indicate the errors in EW measurements, the horizontal the exposure time of the spectra. In the first three nights the variability is of the order of the estimated errors. On 990106 and 990107 it is more or about 5 times the error of the measurements.

night:

$$\sigma = \sqrt{\frac{1}{(N-1)\bar{f}_\lambda^2} \sum_{i=1}^N (f_{i\lambda} - \bar{f}_\lambda)^2}, \quad (1)$$

where N is the number of spectra, $f_{i\lambda}$ is the flux at wavelength λ of the i^{th} spectrum and $\bar{f}_\lambda$ is the flux at that wavelength averaged over all spectra. The σ in the continuum should correspond to $(\text{SNR})^{-1}$, and hence a peak in σ should determine the significance of a variation.

One other statistical technique which is particularly useful to accurately determine the spectral location of variability is the temporal variance spectrum (Prinja et al. 2003; Fullerton, Gies & Bolton 1996):

$$(TVS)_\lambda = \frac{s_0^2}{(N-1)} \sum_{i=1}^N \frac{(f_{i\lambda} - \bar{f}_{w\lambda})^2}{s_i^2 f_{i\lambda}} \quad (2)$$

where $\bar{f}_{w\lambda}$ is the weighted mean of the normalised intensity, s_i is the inverse of the SNR of spectrum i measured in the continuum, and $s_0^2 = [1/N \sum_{i=1}^N s_i^{-2}]^{-1}$.

The results of calculations are presented, together with all spectra, in Fig.3.

As can be seen, the fractional variance spectra do not indicate variability in the first three nights (980415, 980802, and 980803), however the TVS indicates some variability in 980415. A clear peak in both σ and the TVS is visible on 990106 and 990107 at about velocity ≈ 0 km s $^{-1}$.

Following Fullerton, Gies & Bolton (1996), if the TVS for a spectral feature is above a specified level of significance then the null hypothesis of “no variability” can be rejected. The statistical distribution of the TVS is governed by the reduced chi-squared distribution with $N-1$ degrees of freedom and scaled with SNR ($\text{TVS} \sim s_0^2 \chi_{N-1}^2$). For the nights considered here, the statistical significance of 1% corresponds to $\text{TVS}^{1/2} = 3.6-5.0\%$, depending on the number of spectra and SNR. In the central part of the line the TVS is well above this level for 990106 and 990107 (Fig.3). The peaks in σ correspond practically to the same confidence level of 3 standard deviations from the mean. This clearly shows that in these two nights there is statistically significant variability in the H α line profile, at a confidence level of 0.99, at a time resolution of ~ 10 min.

3.2. Variability in January 1999

For the two nights with detected variability, we subtract from the individual spectra the average spectrum, the minimum spectrum and the maximum spectrum. The average spectrum is defined as the average of the normalised flux for every pixel. The minimum (maximum) spectrum is defined as the minimal (maximal) value for each pixel. We do not know *a priori* whether we have variability due to a pure emission, pure absorption, or an admixture of both. The results for 990106 are presented in Fig. 4. As can be seen, variability exists in all three panels. This indicates additional absorption/emission with equivalent width of approximately 1 Å, located in the central part of the line.

To exclude the possibility that the observed variability is an artifact of data reduction we performed tests with data processing and normalisation, which showed that the detected variability on 990106 and 990107 is not an artificial result. Fig.5b shows the results from two different extractions and normalisations which have identical σ peaks.

Although the sky conditions looked like photometric during the time of the observations we can not exclude the possibility that the variable atmospheric lines and small wavelength shifts from spectrum to spectrum can result in spurious variability. To test the variability these effects could produce, we artificially introduced random shifts in wavelength up to 3 km s $^{-1}$ [the biggest shifts we detected were < 0.25 px (2.5 km s $^{-1}$)]. We also isolated atmospheric absorption lines from a high SNR spectrum of Spica (α Vir) and introduced variable atmospheric absorption in the spectra. We find that both effects can produce variability, but this variability is lower than the noise in TVS and σ and can not account for the observed peaks.

It is known (i.e. Zamanov & Bruch 1998) that the hot continuum, which flickers on time scales of minutes with amplitude 0.2-0.3 mag in U, contributes about 15% of the average flux in the V band. To check whether the variability detected in H α is due to the changing continuum level caused by the flickering of the hot component, we simulate σ in such a case by supposing that the hot continuum flickers with amplitude $\pm 25-30\%$ and contributes 15% of the flux in the continuum around H α . The amplitude is chosen to be slightly higher than the highest one observed in the U band after the subtraction of the red giant contribution (see Zamanov & Bruch 1998). To this variable hot continuum we add constant red giant and H α emission. The simulated σ and TVS are shown in Fig.5. The simulated quantities are different from the “observed” ones as (i) they have smaller amplitude and (ii) they are wider and spread over the whole FWZI of the H α emission, while the observed ones are located in the central part of the line only. This indicates that detected variability is not due to the continuum variability but is intrinsic to the line.

4. DISCUSSION

The FWZI of the H α emission line is 800 km s $^{-1}$, with wings of the line formed in the immediate vicinity of the white dwarf (see also Stanishev et al. 2004). However, the detected variability appears in the central part of the line and is confined to $\Delta V \approx \pm 100-120$ km s $^{-1}$ from the systemic velocity. This indicates that this variability originates in the outer parts of the H α emitting region. Assuming Keplerian motion, a 1.4 M $_{\odot}$ white dwarf, and inclination 67 $^{\circ}$ (i.e. Stanishev et al. 2004 and references therein) this corresponds to a distance $\geq 20-25 R_{\odot}$.

Mass accretion in T CrB is occurring via Roche lobe overflow. As the stream of gas flows away from L $_1$ it forms a ring (see Verbunt 1982). The position of the ring approximately defines the outer edge of the accretion disk. The radius of the ring can be estimated from (Hessman & Hopp 1990): $r_r/a = 0.0859(q)^{-0.426}$, where a is the semi-major axis and q is the mass ratio of giant to white dwarf. For T CrB $q = 0.82 \pm 0.10$ and $a \approx 210 R_{\odot}$ (Stanishev et al. 2004), which places the ring at 20 $R_{\odot}$. At this location, the Keplerian velocity is about 115 km s $^{-1}$, which is similar to ΔV of the detected variability. However, the Keplerian time scale is about 18 days, which is not comparable with the timing of our observations. This suggests that the variability is not produced by an inhomogeneous disc structure.

The observed behaviour of H α emission from T CrB with time resolution 10-15 minutes is different from that of CH Cyg and of the variable winds observed in some CVs. We do not see ejected blobs, like those observed in CH Cyg (by Tomov et al. 1996), nor high velocity components observed in the variable disk winds of some CVs (Prinja et al. 2003). It might be that they are not detectable in our H α profiles, even though they may exist and could be detected in the UV, where the hot component is the dominant source of radiation.

One possible explanation for the detected variability could be variable absorption from the wind of the giant or from the accretion stream. It is worth noting, that no variability in the line is observed at orbital phases where the line of sight passes through the expected position of maximum absorption from the red giant wind. Another




FIG. 3.— The $H\alpha$ profiles of T CrB for 5 nights of observations, normalised to the local continuum. The X-axis is in heliocentric velocity. σ and the TVS for all nights are plotted. The dotted horizontal line indicates a 1% threshold for the detection of variability. For first 3 nights σ does not indicate any variability, while the TVS is marginally above the 1% significance level on the night of 980415. On 990106 and 990107 well defined peaks, both in σ and in the TVS, are clearly visible in the central part of the line, corresponding to velocity -100 to +100 km/s. This demonstrates that there is a statistically significant variability in the profile of $H\alpha$ line with time resolution ~ 10 min. The probability that this is a real variability is ≥ 0.99 .

cause could be additional emission from the area where the accretion stream hits the disk (the hot spot). It is possible that a variable component with $EW \leq 0.5 \text{ \AA}$ always exists, but when the line is strong it is lost in the line (e.g. 980415).

An additional plausible explanation is that the variable continuum arising from the central part of the accretion disk (where the flickering in U arises) changes the ionisation state of the outer regions of the (flared) disk, from which the central part of the emission lines arise. This possibility deserves further exploration.

5. CONCLUSIONS

We searched for rapid $H\alpha$ variability in the spectra of the recurrent nova T CrB. On 2 nights out of 5, we detect statistically significant variability at time resolution of 10-20 minutes. The detected variability is confined to the central part of the line profile at $\pm 100 \text{ km s}^{-1}$, which suggests that variations are produced in the outer parts of the accretion disk ($> 20 R_{\odot}$). On these two nights there is evidence for variability of the total $EW(H\alpha)$ (in all cases it is calculated relative to the average value, see Table 1) with amplitude $\pm 8\%$ for 990106 and $\pm 6\%$ for




FIG. 4.— Night 990106: from the individual spectra have been subtracted the average spectrum (left panel), the minimum (middle panel), and the maximum spectrum (right panel). Zero of the velocity axis corresponds to the systemic velocity $\gamma = -27.79$ km s $^{-1}$ (Fekel et al 2000).




FIG. 5.— (a) H α profiles observed on 990106 (the same as in Fig.3). (b) σ for 990106, solid and dashed lines are for two deliberately different extraction and normalisation procedures. (c) simulated σ as a result of hot continuum variability. (d) simulated TVS as a result of hot continuum variability. During the simulation we assumed that the hot continuum contributes 15% of the flux on average and it varies with $\pm 30\%$ (dashed line) and $\pm 25\%$ (solid line). Zero of the velocity axis corresponds to the systemic velocity γ .

990107, which is approximately five times the error of the individual measurements.

For the other 3 nights of observations we do not detect line profile changes. We place upper limits on the variability of the total EW(H α) $\Delta\text{EW}(\text{H}\alpha) \pm 2\%$ for 980415, $\pm 4\%$ for 980802 and $\pm 3\%$ for 980803.

More extensive spectral observations with better time resolution, combined with photometry are necessary to define what influences the H α variability and whether this variability is directly connected with the flickering.

REFERENCES

- Anupama, G. C. & Prabhu, T. P. 1991, MNRAS, 253, 605
- Anupama, G. C. & Mikolajewska, J. 1999, A&A, 344, 177
- Dobrzycka, D., Kenyon, S. J., & Milone, A. A. E. 1996, AJ, 111, 414
- Fekel, F. C., Joyce, R. R., Hinkle, K. H., & Skrutskie, M. F. 2000, AJ, 119, 1375
- Fullerton, A. W., Gies, D. R., & Bolton, C. T. 1996, ApJS, 103, 475
- Hessman, F. V. & Hopp, U. 1990, A&A, 228, 387
- Iijima, T. 1990, Journal of the American Association of Variable Star Observers (JAAVSO), 19, 28
- Kafka, S., Tappert, C., Honeycutt, R. K., & Bianchini, A. 2003, AJ, 126, 1472
- Prinja, R. K., Knigge, C., Ringwald, F. A., & Wade, R. A. 2000a, MNRAS, 318, 368
- Prinja, R. K., Ringwald, F. A., Wade, R. A., & Knigge, C. 2000b, MNRAS, 312, 316
- Prinja, R. K., Long, K. S., Froning, C. S., Knigge, C., Witherick, D. K., Clark, J. S., & Ringwald, F. A. 2003, MNRAS, 340, 551
- Ringwald, F. A. & Naylor, T. 1998, AJ, 115, 286
- Selvelli, P. L., Cassatella, A., & Gilmozzi, R. 1992, ApJ, 393, 289
- Sokoloski, J. L., Bildsten, L., & Ho, W. C. G. 2001, MNRAS, 326, 553
- Sokoloski, J. L., 2002, in press, astro-ph/0209101
- Stanishev, V., Zamanov R., Tomov N., Marziani P., 2004, A&A, 415 609
- Tomov, T., Kolev, D., Munari, U., & Antov, A. 1996, MNRAS, 278, 542
- Tomov, T., Kolev, D., Munari, U., Sostero, G., & Lepardo, A. 1995, A&A, 300, 769
- Zamanov, R. K. & Bruch, A. 1998, A&A, 338, 988
- Zamanov, R. & Marti, J. 2001, IBVS 5013
- Zamanov, R. K., Bode, M.F., Stanishev, V., Martí, J., 2004, MNRAS, in press, astro-ph/0402465
- Verbunt, F. 1982, Space Science Reviews, 32, 379
- Witherick, D. K., Prinja, R. K., Howell, S. B., & Wagner, R. M. 2003, MNRAS, 346, 861
- Warner, B. 1995, Cataclysmic Variable Stars, Cambridge Astrophysics Series, Cambridge, New York: Cambridge University Press, 1995.